

DOCKET NO.: CARP-0057

PATENT

APPENDIX B

Claim Limitation	Support in 1989 GB Application
28. A humanized immunoglobulin having complementarity determining regions (CDRs) from a donor immunoglobulin and heavy and light chain variable region frameworks from human acceptor immunoglobulin heavy and light chains	See page 1, lines 1-2 and 10-20; page 5, lines 8, to page 6, line 4; , and page 8.
which humanized immunoglobulin specifically binds to an antigen with an effective antigen binding affinity	See page 5, lines 1-7; page 22, lines 27-35, page 23, lines 5-9, page 24, lines 1-4; page 25, lines 27-33; page 26 last paragraph.
wherein said humanized immunoglobulin comprises amino acids from the donor immunoglobulin framework outside both the Kabat CDRs and the structural loop CDRs of the variable regions,	See page 5, lines 1-7; page 26, last paragraph, to page 27, top paragraph.
wherein the donor amino acids replace corresponding amino acids in the acceptor immunoglobulin heavy or light chain frameworks,	See page 5, line 8, to page 6, line 4; page 7, lines 5-20.
and each of said donor amino acids contributes to antigen binding as determined by X-ray crystallography.	Page 18, lines 11-17, and lines 33-37, and Figs. 20-21 of the application as filed reference residues that may "contribute to antigen binding" as determined using X-ray crystallography. Residues 48, 49, 71, 73, 76, 78, 88, and 91 are identified in Figure 21.
49. A humanized immunoglobulin having complementarity determining regions (CDRs) from a donor immunoglobulin and heavy and light chain variable region frameworks from human acceptor immunoglobulin heavy and light chains	See page 1, lines 5-16, and page 7, line 32, through page 8, line 21.

DOCKET NO.: CARP-0057

PATENT

which humanized immunoglobulin specifically binds to an antigen with an affinity constant of at least 10^8 M^{-1} ,	See page 11, lines 27-30.
wherein said humanized immunoglobulin comprises amino acids from the donor immunoglobulin framework outside both the Kabat CDRs and the structural loop CDRs of the variable regions,	See page 6, lines 14-23, page 8, lines 13-16, and page 19, line 16, to page 20, line 15.
wherein the donor amino acids replace corresponding amino acids in the acceptor immunoglobulin heavy chain framework	See page 6, line 12, to page 7, line 5.
at residues 48, 49, 71, 73, 76, 78, 88, and 91.	See Light chain 341B of Table 1, page 20.